The 2019 Measles Outbreak



2019's outbreak is the largest since 1994.



This is the greatest number of cases reported in the U.S. since measles was eliminated in 2000.

Complications of Measles

- Pneumonia is the #1 cause of pediatric measles-related deaths
- Diarrhea, stomatitis, conjunctivitis, and keratitis (may lead to blindness) may occur
- Encephalitis is rare


What are the symptoms?

- Red maculopapular rash that starts on head and spreads down
- Becomes confluent
- Fades in order of appearance

Remember the 3 Cs:

- Cough
- Coryza (runny nose)
- Conjunctivitis (red eyes)





First dose at 12-15 months 95% protective Second dose at 4-6 years 99% protective



Testing for Suspected Measles

- Consult with the local/state health departments and local Infectious Diseases specialists for guidance
- Measles IgM (may need to be repeated if initial test is negative because false-negatives may be noted during the first 72 hours of rash onset)
- Serum and throat swabs for measles
 RNA by PCR (send out test ran at state public health laboratories)
- Measles IgG (presence suggests immunity)



To learn more about vaccinations or to schedule a Maximizing Office Based Immunization (MOBI) training with your practice, please contact Lory Sheeran Winland at **Iwinland@ohioaap.org**.

American Academy of Pediatrics (dedicated to the health of all children^{**}

